

**Atrial Fibrillation Association** (AFA) is a registered charity.

Internationally, AFA works closely with patients, carers, medical professionals, service providers, service payers affiliated groups and allied professionals to:

- provide support and information
- advance the education of the medical professionals
- raise awareness amongst the general public
- promote research into the management of Atrial Fibrillation

In Wales, many thousands of preventable strokes occur every year leading to thousands of early deaths and a devastating burden on individuals, families and society in terms of disability, medical and social care costs, and loss of working hours and tax revenues.

Atrial fibrillation (AF) is a common heart rhythm disorder associated with deadly and debilitating consequences including heart failure, stroke, poor mental health, reduced quality of life and death.<sup>i</sup>

### Key points of AFA evidence

1. AF patients suffer a disproportionate number of strokes which are, in turn, disproportionately fatal, debilitating, expensive and likely to recur.
2. AF detection and diagnosis is low, leaving an estimated 50% of patients undiagnosed. **Opportunistic screening**, has been shown to be both effective and cost efficient at finding AF patients, however it is not widely used.
3. **Guidance and guideline adherence** is poor and consequently leaves AF patients at risk of avoidable strokes.
4. The main current therapy option, warfarin, which can effectively prevent many of these strokes is actually effective in only 18-21% of AF patients, due to under prescribing especially in those at high risk of stroke.
5. Too often, those at most risk, frequently the elderly, are prescribed aspirin, which only reduces the risk of stroke by 22% and increases their risk of a bleed to equal that of warfarin.
6. For those who often fall out of therapeutic levels (> 60% outside of target therapeutic range), warfarin is of little or no benefit.
7. The Inquiry Committee may wish to consider seeking evidence from all relevant professionals, including heart rhythm specialists.

### Supporting evidence

AFA is mindful that budgetary pressures are ever-present and inevitable, and as a result, cost effectiveness has to be a reasonable expectation when comparing guidance it is important to consider both cost and effectiveness. This difference is probably best summarised but the QIPP, Right Care programme, 'Commissioning for Value':

*'...value must also be measured by outputs, not inputs. Hence it is patient health results that matter...'*

The AFA has amassed and documented many thousands of experiences that have been shared with us by patients, their caregivers and health care providers. These accounts are an excellent representation of the "health results" of patients suffering from AF in the UK today. In light of this amassed patient feedback and respected published data we have formulated the following response on behalf of patients suffering from AF.

**1. AF is the most powerful independent risk factor for stroke and results in strokes that are more severe, more likely to disable, more likely to kill and more likely to recur.**

Atrial Fibrillation (AF) is a common heart rhythm disorder associated with deadly and debilitating consequences including heart failure, stroke, poor mental health, reduced quality of life and death. [26] Today, more than 51,000 of the Welsh population have been diagnosed with AF, [28] yet experts suggest that up to half of all AF patients have not yet been detected. Among many damaging and debilitating consequences, AF increases an individual's risk of suffering a stroke by five times. [31] This effect alone results in considerable disability and death, [27,32] not to mention avoidable millions in healthcare expenditure [28] that NHS Wales cannot afford. For example, patients with primary or secondary diagnosis of AF occupied almost 308,000 bed days in 2008, at a cost to NHS Wales of £100 million [265]. Strokes kill about 1325 people in Wales each year. AF is known to be responsible for almost one quarter of these strokes, the health and social care costs of AF-related strokes in Wales are expected to reach £46.3 million per year [266]. These figures exclude the economic burden on NHS Wales and the Welsh economy.

Atrial Fibrillation is known to be responsible for 45% of all embolic strokes, resulting in more than 12,500 strokes per year in England and Wales. The strokes suffered by people with AF are also more severe, [82] they are more frequently fatal [83 84] and they are more likely to lead to disability, [83 85 86 82] increased healthcare costs [89] and extended hospital care than strokes in patients without AF. [82] Moreover, AF-related strokes are more likely to happen again [89], adding not just to the risk of future strokes, but also to the potential for increased patient anxiety and a further reduction in quality of life. AF-related strokes kill nearly twice as frequently as non-AF strokes. [82,83,84]

The medical cost of a stroke in first year is £9,500 - £14,000 per stroke. Embolic, and hence AF related, strokes are likely to be represented at the high end of this range. [93,98,100,103,104] These costs do not include continuing costs after first year, nor do they include costs associated with long term disability or the human cost, which is incalculable.

**2. AF detection and diagnosis is low, leaving an estimated 50% of patients undiagnosed**

Without effective detection and diagnosis of AF up to half the patients affected will never be identified. If a lack of detection and diagnosis continues, then many patients will be denied the opportunity to benefit from treatments that can dramatically reduce their risk of stroke. Existing research suggests that routine pulse screening has a role to play, as does public education on the need to investigate an irregular pulse.

**Opportunistic screening, has been shown to be both effective and cost efficient at finding AF patients, however it is not widely used.**

The role and value of screening programmes following the positive results of the SAFE study [266] illustrate the value of this in both finding patients with AF and reducing stroke risk, and ultimately stroke events. This has been more recently shown in Wrexham [268] where The Wrexham primary care AF model pilot launched on 1st May 2007 in four general practice surgeries under the Local Health Boards and ran for six months with positive results and findings. The audit of the pilot evaluated the view of existing patients on the AF registers in the surgeries as well as those patients that were identified through the routine screening using the manual pulse check. Seven new AF patients were found during opportunistic checks as part of the pilot and 68 patients were found to be on inappropriate or no thromboprophylaxis, which prompted further review by the GPs. Four

years since the start of the pilot the team are still reaping the benefits of the work undertaken in primary care.

NICE also recommends that all patients with an irregular pulse receive an ECG to make a diagnosis. However, The SAFE study found that GP and practice nurse performance in interpreting ECGs was not encouraging, identifying another potential challenge to the effective diagnosis of AF in primary care.

**The need for routine, opportunistic screening, especially amongst those at greater risk of developing AF is essential if lives are to be saved and costs due to stroke events, reduced.**

### 3. Guidance adherence is poor and consequently leaves AF patients at risk of avoidable strokes

There remains poor adherence to authoritative and national guidelines. Even NICE's own data shows that of all those with AF who should be on warfarin, almost half are not. Yet, when asked, physicians demonstrate both awareness of the guidelines and agreement with them. One study documented the medications being taken by AF patients when they suffered an ischemic stroke. It found that only 10% of these patients had been taking an effective dose of an anticoagulant. Nearly a third were on no antithrombotic treatment at all (29%). A further 29% were on aspirin and another 29% were on a non-therapeutic dose of warfarin.




Figure 10. Medications received before admission to hospital by patients with known atrial fibrillation who suffered an acute ischaemic stroke: only 10% of patients had received warfarin at a therapeutic dose. Adapted with permission from Gladstone

Adherence to Guidelines for the prevention of stroke in patients with AF may be low for several reasons. These include difficulties in maintaining INR within the therapeutic range and physicians' concerns about bleeding risk, particularly in the elderly. [35]

Quality Outcomes Framework (QOF) was designed to reward GPs for the quality of the care that they provided, instead of for how many patients they treated. Currently, GPs can work to secure up to 1,000 QOF points by meeting predetermined performance targets in four broad areas: Clinical, Organisational, Patient Experience, and Additional Services. A total haul of 1,000 points represents an additional payment to GPs of over £13,000.

Specifically for AF, within the current QOF scheme, up to 12 points are available for GPs achieving a high percentage of 'patients with atrial fibrillation who are currently treated with anticoagulation drug therapy or antiplatelet therapy.'

It is reasonable to expect that many patients diagnosed with AF will already be taking aspirin for another condition. It is also relatively simple to start and manage a patient on aspirin (antiplatelet

therapy) compared to warfarin (anticoagulation therapy). Consequently, the way the target is written enables GPs to receive the maximum QOF reward just by having AF patients on aspirin, even if none of them is on warfarin.

Consequently, QOF today provides virtually no incentive for GPs to put patients on warfarin in accordance with the NICE 2006 or the ESC 2010 guidelines.

AFA would suggest that guidelines and rewards encourage

- An increase in the percentage of patients with Atrial Fibrillation in whom stroke risk has been assessed using the CHADS<sub>2</sub> risk stratification scoring system in the previous 15 months
- In those patients with Atrial Fibrillation in whom there is a record of a CHADS<sub>2</sub> score of  $\geq 1$ , an increase the percentage of patients who are receiving anticoagulants

**Adherence to agreed guidance is essential if AF patients are to be properly assessed and treated to reduce stroke.**

#### 4. Aversion to warfarin leaves thousands of patients at unnecessarily high risk of stroke

In clinical trials warfarin has been associated with a stroke risk reduction in AF patients of 50%-70%. However, this potential is not being realised in routine clinical practice, leaving thousands at risk of preventable strokes.

Warfarin is currently recommended in UK and European guidelines as first-line therapy in patients with AF and a moderate or high risk of developing stroke. [138 139] Despite evidence that following the guidelines results in improved patient outcomes, [166] there is significant under-use of warfarin. Thus, many patients with AF and a moderate-to- high risk of stroke do not receive anticoagulant therapy and therefore remain at high risk for stroke. [156 153]

NICE data shows that of all those with AF who should be on warfarin, almost half are not. [179]. In a study conducted in seven European countries, it was found that only 8.4% of patients with AF who had a stroke were receiving anticoagulants at the time of their stroke, and the proportion decreased by 4% per year with increasing age. [181] A review of the scientific literature from 2000 indicated that only 15–44% of eligible patients with AF were receiving warfarin.[180] Yet, when asked, physicians demonstrate both awareness of the guidelines and agreement with them, despite not treating patient in accordance with those guidelines. [182] This further highlights the discrepancy that is often found between trial results and what happens in clinical practice. Another study documented the medications being taken by AF patients when they suffered an ischemic stroke. It found that only 10% of these patients had been taking an effective dose of an anticoagulant. Nearly a third were on no antithrombotic treatment at all (29%). A further 29% were on aspirin and another 29% were on a non-therapeutic dose of warfarin. According to recent surveys in different parts of Europe, the proportion of patients with AF at high risk of stroke who are receiving adequate anticoagulation is most commonly around 54–61% [185,172]

Warfarin is under-prescribed for many reasons including the complexity of dosing and patient management as well as fear of the associated bleeding risks. Consequently, almost half the AF patients for whom warfarin is indicated are not on warfarin and remain at extremely high risk of severe, debilitating and expensive strokes.

Management of warfarin is complex and time-consuming for primary care physicians. It is also recognised that those at greatest risk, the elderly, are less likely to be given warfarin because of perceived fear of complications.

In centres where clinicians with a special interest in AF have set up clinics and outreach support, such as Prince Philip Hospital, Llanelli, commitment of a truly multidisciplinary team, has led to a

successful "one-stop service" service for those referred with AF. When started, only 22% of the patients referred to Dr Izzat with a CHADS<sub>2</sub> of more than 1 were adequately anti-coagulated with warfarin, this has now improved to just over 40% and plans to improve this further are in hand. Knowledge of and adherence to current guidance, effective use of validated risk assessment schema and appropriate use of anticoagulation has lead to successful stroke prevention – and with no extra finance.

**5. Too often, those at most risk, frequently the elderly, are prescribed aspirin, which only reduces the risk of stroke by 22% and increases their risk of bleed to equal that of warfarin.**

Many physicians resist the use of warfarin in the elderly, largely on grounds of safety. Research has demonstrated repeatedly that physicians over-estimate the risk of bleeding associated with the use of warfarin and under-estimate its benefits in preventing thromboembolism and stroke; conversely, they have been shown to under-estimate the bleeding risk of aspirin therapy and over-estimate its benefits. [196,188,201] As a result, eligible patients are not receiving therapy that could prevent strokes. [18] For many physicians, bleeding risk is a particular concern in the elderly, who are more prone to falls, more likely to have suffered previous major bleeds, and who are subject to many additional problematic factors associated with old age. [202,204] While the bleeding risk with warfarin is no worse than that with aspirin, physician experience of major bleeding events associated with warfarin can profoundly reduce prescription of warfarin. [205] A study investigated the behaviour of physicians treating AF patients who had bleeds while on warfarin. Patients treated in the 90 days after the physician had encountered a bleeding event were significantly less likely to receive a prescription for warfarin than patients treated before the bleed. [205] In contrast, having a patient who experienced a Stroke while not receiving warfarin did not influence prescribing behaviour with subsequent patients. [205] In other words, a bleeding event may make a physician less likely to prescribe an anticoagulant but a stroke does not increase the likelihood that a physician will prescribe an anticoagulant. There are large numbers of younger patients who according to guidance should be prescribed anticoagulants including 'those with a history of stroke and those aged 65 years or over with one of the following: diabetes, coronary artery disease, or hypertension', but who simply are not receiving it, whether this be through their choice or physician assessment. **AFA would propose that guidelines recommending that patients be assessed for risk using the CHADS<sub>2</sub> and the CHADS<sub>2</sub>VASc<sub>2</sub> system should be adopted.**

**6. For those patients on warfarin large numbers of patients are difficult to control and spend >60% outside the target therapeutic range – rendering warfarin of no benefit.**

Research has shown that AF patients in routine clinical care were able to maintain a target INR for over half the time (56%). Of the considerable remaining time, patients were above the target range for 30%, and below the target range for 14%. [184]. If around half of all patients in need of anticoagulation aren't prescribed warfarin [179] and if those who are have either ineffective or unsafe blood levels of warfarin for nearly half of the time, [184] then perhaps only a quarter of patients at any one time receive the therapy they need to safely lower their risk of stroke. This becomes ever more worrisome when considering experts' estimates that only about half of all AF patients are actually diagnosed. The vast majority of these undiagnosed patients would be expected to be at moderate or high risk of stroke, [190] and, hence, in need of warfarin therapy according to

the ESC 2010 guidelines. Yet perhaps only a fifth of patients in need of warfarin to reduce risk of stroke are actually receiving safe and effective anticoagulation treatment at any time.

-Effective monitoring is essential if at risk AF patients are to be protected.

-An education programme, in line with recommended guidance is required to increase awareness and understanding of the importance of appropriate anticoagulation therapy in AF patients.

**7. The Inquiry Committee may wish to consider seeking evidence from all relevant professionals, including heart rhythm arrhythmia specialists.**

AF is largely managed in primary care, but cardiologist who are arrhythmia specialists also play a key role in local training, referrals and care of more challenging AF patients. Cardiac Networks and cardiology arrhythmia specialists are also instrumental in informing and supporting local guideline adherence in order to PREVENT stroke in AF patients.

AFA has worked with the Cardiac Networks and Arrhythmia specialist in Wales, and suggests that the Inquiry would benefit from seeking expert advice from these sources. AFA would be very happy to highlight Welsh centres where specialist clinicians are working to reduce both the burden of AF and the risk of stroke amongst the AF population.

## References

- 1 Office of National Statistics Health Statistics Quarterly (12) Winter 2001 "Stroke incidence and risk factors in a population based cohort study"; Scottish Stroke Care Audit 2005/2006.
- 2 Adamson, J., Beswick, A. and Ebrahim, S., "Stroke and Disability" in Journal of Stroke and Cerebrovascular Diseases, Vol 13, No 4, 2004
- 3 Reducing Brain Damage: Faster access to better stroke care, National Audit Office Report. Department of Health, 2005
- 4 Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. Stroke 1991;22:983–8
- 5 Marini C, De Santis F, Sacco S et al. Contribution of atrial fibrillation to incidence and outcome of ischemic stroke: results from a population- based study. Stroke 2005;36:1115–19
- 6 Lamassa M, Di Carlo A, Pracucci G et al. Characteristics, outcome, and care of stroke associated with atrial fibrillation in Europe: data from a multicenter multinational hospital- based registry (The European Community Stroke Project). Stroke 2001;32:392–8
- 7 Wolfe CD. The impact of stroke. Br Med Bull 2000;56:275–86
- 8 White CL, Poissant L, Cote-LeBlanc G et al. Long-term caregiving after stroke: the impact on caregivers' quality of life. J Neurosci Nurs 2006;38:354–60
- 9 Fuster V, Ryden LE, Cannom DS et al. ACC/AHA/ESC 2006 Guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 Guidelines for the Management of Patients With Atrial Fibrillation): developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. Circulation 2006;114:e257–354
- 10 Kirchhof P, Auricchio A, Bax J et al. Outcome parameters for trials in atrial fibrillation: recommendations from a consensus conference organized by the German Atrial Fibrillation Competence NETWORK and the European Heart Rhythm Association. Europace 2007;9:1006–23
- 11 Lip GYH et al. Stroke 2002;33:238–44
- 12 Go AS, Hylek EM, Phillips KA et al. Prevalence of diagnosed atrial fibrillation in adults: national implications for rhythm management and stroke prevention: the AnTicoagulation and Risk Factors in Atrial Fibrillation (ATRIA) Study. JAMA 2001;285:2370–5
- 13 Miyasaka Y, Barnes ME, Gersh BJ et al. Secular trends in incidence of atrial fibrillation in Olmsted County, Minnesota, 1980 to 2000, and implications on the projections for future prevalence. Circulation 2006;114:119–25
- 14 Gage BF, Waterman AD, Shannon W et al. Validation of clinical classification schemes for predicting stroke: results from the National Registry of Atrial Fibrillation. JAMA 2001;285:2864–70
- 15 Briffa T, Hickling S, Knuijan M et al. Long term survival after evidence based treatment of acute myocardial infarction and revascularisation: follow-up of population based Perth MONICA cohort, 1984–2005. BMJ 2009;338:b36
- 16 Lip GY, Lim HS. Atrial fibrillation and stroke prevention. Lancet Neurol 2007;6:981–93
- 17 Wittkowsky AK. Effective anticoagulation therapy: defining the gap between clinical studies and clinical practice. Am J Manag Care 2004;10:S297–306
- 18 Gladstone DJ, Bui E, Fang J et al. Potentially preventable strokes in high-risk patients with atrial fibrillation who are not adequately anticoagulated. Stroke 2009;40:235–40
- 19 Hirsh J, Dalen J, Anderson DR et al. Oral anticoagulants: mechanism of action, clinical effectiveness, and optimal therapeutic range. Chest 2001;119:8S–21S
- 20 Turpie AG, Warfarin replacements: Mechanisms underlying emerging agents. Can J Cardiol 2008;24 Suppl C:56–60C
- 21 Frykman V, Beerman B, Ryden L et al. Management of atrial fibrillation: discrepancy between guideline recommendations and actual practice exposes patients to risk for complications. Eur Heart J 2001;22:1954–9
- 22 Man-Son-Hing M, Laupacis A. Anticoagulant-related bleeding in older persons with atrial fibrillation: physicians' fears often unfounded. Arch Intern Med 2003;163:1580–6
- 23 Lip GY, Zarifis J, Watson RD et al. Physician variation in the management of patients with atrial fibrillation. Heart 1996;75:200–5
- 24 Daniel K, Wolfe CD, Busch MA et al. What are the social consequences of stroke for working-aged adults? A systematic review. Stroke 2009;40: e431–40
- 25 Bungard TJ, Ghali WA, Teo KK et al. Why do patients with atrial fibrillation not receive warfarin? Arch Intern Med 2000;160:41–6
- 26 Stewart S, Hart CL, Hole DJ et al. A population-based study of the long- term risks associated with atrial fibrillation: 20-year follow-up of the Renfrew/Paisley study. Am J Med 2002;113:359–64
- 27 Kannel WB, Wolf PA, Benjamin EJ et al. Prevalence, incidence, prognosis, and predisposing conditions for atrial fibrillation: population-based estimates. Am J Cardiol 1998; 82:2N–9N
- 28 The Office of Health Economics Estimating the direct costs of atrial fibrillation to the NHS in the constituent countries of the UK and at SHA level in England, 2008 November 2009. London
- 29 Lloyd-Jones DM, Wang TJ, Leip EP et al. Lifetime risk for development of atrial fibrillation: the Framingham Heart Study. Circulation 2004;110: 1042–6
- 30 Eric J, Feuer EJ, Wun LM, Boring CC et al. The Lifetime Risk of Developing Breast Cancer. J Natl Cancer Inst (1993) 85 (11): 892-897
- 31 Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. Stroke 1991;22:983–8
- 32 Marini C, De Santis F, Sacco S et al. Contribution of atrial fibrillation to incidence and outcome of ischemic stroke: results from a population- based study. Stroke 2005;36:1115–19
- 33 Friberg J, Buch P, Scharling H, et al. Rising rates of hospital admissions for atrial fibrillation. Epidemiology 2003;14:666–72.
- 34 Miyasaka Y, Barnes ME, Gersh BJ et al. Secular trends in incidence of atrial fibrillation in Olmsted County, Minnesota, 1980 to 2000, and implications on the projections for future prevalence. Circulation 2006;114:119–25
- 35 Banach M, Mariscalco G, Urgulan M et al. The significance of preoperative atrial fibrillation in patients undergoing cardiac surgery: preoperative atrial fibrillation – still underestimated opponent. Europace. 2009; 10:1266-70
- 36 McCabe PJ, Schumacher K, Barnason SA. Living with atrial fibrillation: a qualitative study. J Cardiovasc Nurs. 2011 Jul-Aug;26(4):336-44.
- 37 Thrall G, Lip GY, Carroll D, Lane D. Depression, anxiety, and quality of life in patients with atrial fibrillation. Chest. 2007 Oct;132(4):1259-64. Epub 2007 Jul 23.
- 38 Fuster V, Ryden LE, Cannom DS et al. ACC/AHA/ESC 2006 Guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 Guidelines for the Management of Patients With Atrial Fibrillation): developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. Circulation 2006;114:e257–354
- 39 Royal College of Physicians. Atrial fibrillation. National clinical guideline for management in primary and secondary care. National Collaborating Centre for Chronic Conditions. <http://www.nice.org.uk/nicemedia/pdf/cg036fullguideline.pdf>. Accessed March 2009
- 40 AF AWARE. AF AWARE cardiology groups call for greater awareness and better education on atrial fibrillation. Press release 2009. <http://www.world-heart-federation.org/press/press-releases/news-details/article/af-aware-cardiology-groups-call-for-greater-awareness-and-better-education-on-atrial-fibrillation/>. Accessed 16 June 2009
- 41 Lip GYH, Kamath S, Jafriet M et al. Ethnic Differences in Patient Perceptions of Atrial Fibrillation and Anticoagulation Therapy. Stroke 2002;33:238–44
- 42 National Institute for Health and Clinical Excellence. Guideline 36 – Atrial fibrillation: the management of atrial fibrillation. <http://www.nice.org.uk/nicemedia/pdf/CG036niceguideline.pdf>. Accessed March 2009
- 43 Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation: a major contrib- utor to stroke in the elderly. The Framingham Study. Arch Intern Med 1987;147:1561–4
- 44 Jorgensen HS, Nakayama H, Reith J et al. Acute stroke with atrial fibrillation. The Copenhagen Stroke Study. Stroke 1996;27:1765–9
- 45 National Institute for Health and Clinical Excellence. Understanding NICE guidance: Atrial fibrillation. <http://www.nice.org.uk/nicemedia/pdf/CG036publicinfo.pdf>. Accessed May 2009
- 46 1. American Heart Association. Heart Disease and Stroke Statistics: 2004 Update. Dallas, Tex: American Heart Association; 2003.
- 47 Murphy NF, Simpson CR, Jhund PS et al. A national survey of the prevalence, incidence, primary care burden and treatment of atrial fibrillation in Scotland. Heart 2007;93:606–12
- 48 Kirchhof P, Bax J, Blomstrom- Lundquist C et al. Early and comprehensive management of atrial fibrillation: executive summary of the proceedings from the 2nd AFNET- EHRA consensus conference „research perspectives in AF“. Eur Heart J 2009;doi:10.1093/eurheartj/ehp235
- 49 Lip GY, Beevers DG, Singh SP et al. ABC of atrial fibrillation. Aetiology, pathophysiology, and clinical features. BMJ 1995;311:1425–8
- 50 Iqbal MB, Taneja AK, Lip GY et al. Recent developments in atrial fibrillation. BMJ 2005;330:238–43
- 51 Gudbjartsson DF, Arnar DO, Helgadóttir A et al. Variants conferring risk of atrial fibrillation on chromosome 4q25. Nature 2007;448:353–7
- 52 Aizer A, Gaziano JM, Cook NR et al. Relation of vigorous exercise to risk of atrial fibrillation. Am J Cardiol 2009; 103:1572–7
- 53 Farrar MW, Bogart DB, Chapman SS et al. Atrial fibrillation in athletes. Mo Med 2006;103:297–301
- 54 A Ruizgomez et al. Incidence of chronic atrial fibrillation in general practice and its treatment pattern. Journal of Clinical Epidemiology 2002 55: 358-363
- 55 Wolfe CD, Rudd AG. The Burden of Stroke White Paper: Raising awareness of the global toll of stroke-related disability and death. <http://www.safestroke.org/Portals/10/FINAL%20Burden%20of%20Stroke.pdf>. Accessed March 2009
- 56 Adamson J, Beswick A, Ebrahim S. Is stroke the most common cause of disability? Journal of Stroke and Cerebrovascular Diseases, 2004 Jul-Aug;13(4):171-7

57 Reducing Brain Damage: Faster access to better stroke care, National Audit Office Report. Department of Health, 2005

58 Adamson, J., Beswick, A. and Ebrahim, S., "Stroke and Disability" in Journal of Stroke and Cerebrovascular Diseases, Vol 13, No 4, 2004

59 Reducing Brain Damage: Faster access to better stroke care, National Audit Office Report. Department of Health, 2005

60 Easton JD, Saver JL, Albers GW et al. Definition and evaluation of transient ischemic attack: a scientific statement for healthcare professionals from the American Heart Association/American Stroke Association Stroke Council; Council on Cardiovascular Surgery and Anesthesia; Council on Cardiovascular Radiology and Intervention; Council on Cardiovascular Nursing; and the Interdisciplinary Council on Peripheral Vascular Disease. Stroke 2009;40:2276–93

61 Office of National Statistics Health Statistics Quarterly (12) Winter 2001 "Stroke incidence and risk factors in a population based cohort study"; Scottish Stroke Care Audit 2005/2006.

62 Stroke Statistics 2009. British Heart Foundation and The Stroke Association.

63 Wolfe, C "The Burden of Stroke" in Wolfe, C, Rudd, T and Beech, R (eds) Stroke Services and Research (1996) The Stroke Association

64 2005 Coronary Heart Disease Statistics. British Heart Foundation

65 World Health Organization. The global burden of disease: 2004 update. [http://www.who.int/healthinfo/global\\_burden\\_disease/2004\\_report\\_update/en/index.html](http://www.who.int/healthinfo/global_burden_disease/2004_report_update/en/index.html). Accessed March 2009

66 Boyle P, Ferlay J. Cancer incidence and mortality in Europe, 2004. Ann Oncol 2005;16:481–8

67 Truelsen T, Piechowski-Jozwiak B, Bonita R et al. Stroke incidence and prevalence in Europe: a review of available data. Eur J Neurol 2006;13:581–98

68 Daniel K, Wolfe CD, Busch MA et al. What are the social consequences of stroke for working-aged adults? A systematic review. Stroke 2009;40: e431–40

69 Gunaratne A, Patel JV, Gammon B et al. Ischemic stroke in South Asians. A review of the epidemiology, pathophysiology, and ethnicity-related clinical features. Stroke 2009;40:e415–23

70 Mayo NE, Wood-Dauphinee S, Ahmed S et al. Disablement following stroke. Disabil Rehabil 1999;21: 258–68

71 Kappelle LJ, Adams HP Jr, Heffner ML et al. Prognosis of young adults with ischemic stroke. A long-term follow-up study assessing recurrent vascular events and functional outcome in the Iowa Registry of Stroke in Young Adults. Stroke 1994;25:1360–5

72 Spielier JF, Lanoe JL, Amarenco P. Socioeconomic aspects of postacute care for patients with brain infarction in France. Cerebrovasc Dis 2002;13: 132–41

73 Bruggenjurgen B, Rossnagel K, Roll S et al. The impact of atrial fibrillation on the cost of stroke: the Berlin acute stroke study. Value Health 2007;10: 137–43

74 Reducing Brain Damage: Faster access to better stroke care. National Audit Office 2005

75 Reducing Brain Damage: Faster access to better stroke care. National Audit Office 2005

76 Reducing Brain Damage: Faster access to better stroke care, National Audit Office Report. Department of Health, 2005.

77 Allender S, Scarborough P, Peto V et al. European Cardiovascular Disease Statistics 2008 Edition. <http://www.heartstats.org/uploads/documents%5Cproof30NOV2007.pdf>. Accessed March 2009

78 Chandratheva A, Mehta Z, Geraghty OC et al. Population-based study of risk and predictors of stroke in the first few hours after a TIA. Neurology 2009;72:1941–7

79 Death rates from stroke, adults aged 65 to 74, 1969 to 2006, England. <http://www.heartstats.org/temp/Figsp1.1f1spweb08.xls>. Accessed March 2009

80 Asplund K, Marké L-A, Terént A et al. Costs and gains in stroke prevention: European perspective. Cerebrovasc Dis 1993;3 (Suppl 1):34–42

81 Lightowers S, McGuire A. Cost-effectiveness of anticoagulation in nonrheumatic atrial fibrillation in the primary prevention of ischemic stroke. Stroke 1998;29:1827–32

82 The Copenhagen Stroke Study. Stroke 1996;27:1765–9

83 Marini C, De Santis F, Sacco S et al. Contribution of atrial fibrillation to incidence and outcome of ischemic stroke: results from a population-based study. Stroke 2005;36:1115–19

84 Steger C, Pratter A, Martinek-Bregel M et al. Stroke patients with atrial fibrillation have a worse prognosis than patients without: data from the Austrian Stroke registry. Eur Heart J 2004;25:1734–40

85 Lamassa M, Di Carlo A, Pracucci G et al. Characteristics, outcome, and care of stroke associated with atrial fibrillation in Europe: data from a multicenter multinational hospital-based registry (The European Community Stroke Project). Stroke 2001;32:392–8

86 Iqbal MB, Taneja AK, Lip GY et al. Recent developments in atrial fibrillation. BMJ 2005;330:238–43

87 Go AS, Hylek EM, Phillips KA et al. Prevalence of diagnosed atrial fibrillation in adults: national implications for rhythm management and stroke prevention: the AnTicoagulation and Risk Factors in Atrial Fibrillation (ATRIA) Study. JAMA 2001;285:2370–5

88 Wattigney WA, Mensah GA, Croft JB. Increased atrial fibrillation mortality: United States, 1980–1998. Am J Epidemiol 2002;155:819–26

89 Winter Y, Wolfram C, Schaeg M et al. Evaluation of costs and outcome in cardioembolic stroke or TIA. J Neurol 2009;256:954–63

90 Schneck M, Lei X. Cardioembolic stroke. eMedicine Neurology 2008. <http://emedicine.medscape.com/article/1160370-overview>. Accessed March 2009

91 Murphy R, Sackley CM, Miller P et al. Effect of experience of severe stroke on subjective valuations of quality of life after stroke. J Neurol Neurosurg Psychiatry 2001;70:679–81

92 Gage BF, Cardinalli AB, Owens DK. The effect of stroke and stroke prophylaxis with aspirin or warfarin on quality of life. Arch Intern Med 1996;156:1829–36

93 Jorgensen HS, Nakayama H, Reith J et al. Acute stroke with atrial fibrillation. The Copenhagen Stroke Study. Stroke 1996;27:1765–9

94 Steger C, Pratter A, Martinek-Bregel M et al. Stroke patients with atrial fibrillation have a worse prognosis than patients without: data from the Austrian Stroke registry. Eur Heart J 2004;25:1734–40

95 Wolfe CD, Rudd AG. The Burden of Stroke White Paper: Raising awareness of the global toll of stroke-related disability and death. <http://www.safestroke.org/Portals/10/FINAL%20Burden%20of%20Stroke.pdf>. Accessed March 2009

96 Lamassa M, Di Carlo A, Pracucci G et al. Characteristics, outcome, and care of stroke associated with atrial fibrillation in Europe: data from a multicenter multinational hospital-based registry (The European Community Stroke Project). Stroke 2001;32:392–8

97 Allender S, Scarborough P, Peto V et al. European Cardiovascular Disease Statistics 2008 Edition. <http://www.heartstats.org/uploads/documents%5Cproof30NOV2007.pdf>. Accessed March 2009

98 Grant JS, Glandon GL, Elliott TR et al. Caregiving problems and feelings experienced by family caregivers of stroke survivors the first month after discharge. Int J Rehabil Res 2004; 27:105–11

99 Young AJ, Rogers A, Addington-Hall JM. The quality and adequacy of care received at home in the last 3 months of life by people who died following a stroke: a retrospective survey of surviving family and friends using the Views of Informal Carers Evaluation of Services questionnaire. Health Soc Care Community 2008;16:419–28

100 Winter Y, Wolfram C, Schaeg M et al. Evaluation of costs and outcome in cardioembolic stroke or TIA. J Neurol 2009;256:954–63

101 Ferro JM. Cardioembolic stroke: an update. Lancet Neurol 2003;2:177–88

102 The Office of Health Economics Estimating the direct costs of atrial fibrillation to the NHS in the constituent countries of the UK and at SHA level in England, 2008 November 2009, London

103 Bruggenjurgen B, Rossnagel K, Roll S et al. The impact of atrial fibrillation on the cost of stroke: the Berlin acute stroke study. Value Health 2007;10: 137–43

104 Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation: a major contributor to stroke in the elderly. The Framingham Study. Arch Intern Med 1987;147:1561–4

105 Fuster V, Ryden LE, Cannom DS et al. ACC/AHA/ESC 2006 Guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 Guidelines for the Management of Patients With Atrial Fibrillation): developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. Circulation 2006;114:e257–354

106 Royal College of Physicians. Atrial fibrillation. National clinical guideline for management in primary and secondary care. National Collaborating Centre for Chronic Conditions. <http://www.nice.org.uk/nicemedia/pdf/cg036fullguideline.pdf>. Accessed March 2009

107 Hobbs FD, Fitzmaurice DA, Mant J et al. A randomised controlled trial and cost-effectiveness study of systematic screening (targeted and total population screening) versus routine practice for the detection of atrial fibrillation in people aged 65 and over. The SAFE study. Health Technol Assess 2005;9:1–74

108 The Task Force for the Management of Atrial Fibrillation of the European Society of Cardiology. Guidelines for the management of atrial fibrillation. European Heart Journal (2010) 31, 2369–2429

109 Aspirin for the prevention of cardiovascular disease: U.S. Preventive Services Task Force recommendation statement. Ann Intern Med 2009;150:396–404

110 Palikhe NS, Kim SH, Park HS. What do we know about the genetics of aspirin intolerance? J Clin Pharm Ther 2008; 33:465–72

111 Murphy NF, Simpson CR, Jhund PS et al. A national survey of the prevalence, incidence, primary care burden and treatment of atrial fibrillation in Scotland. Heart 2007;93:606–12

112 Frost L, Vukelic Andersen L, Godtfredsen J et al. Age and risk of stroke in atrial fibrillation: evidence for guidelines? Neuroepidemiology 2007;28:109–15

113 Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. Stroke 1991;22:983–8

114 Truelsen T, Piechowski-Jozwiak B, Bonita R et al. Stroke incidence and prevalence in Europe: a review of available data. Eur J Neurol 2006;13:581–98

115 Cabin HS, Clubb KS, Hall C et al. Risk for systemic embolization of atrial fibrillation without mitral stenosis. Am J Cardiol 1990;65:1112–16

116 Stewart S, Hart CL, Hole DJ et al. A population-based study of the long-term risks associated with atrial fibrillation: 20-year follow-up of the Renfrew/Paisley study. Am J Med 2002;113:359–64

117 Hughes M, Lip GY. Stroke and thromboembolism in atrial fibrillation: a systematic review of stroke risk factors, risk stratification schema and cost effectiveness data. Thromb Haemost 2008;99:295–304

118 Jorgensen HS, Nakayama H, Reith J et al. Acute stroke with atrial fibrillation. The Copenhagen Stroke Study. Stroke 1996;27:1765–9

119 Lip GY, Lim HS. Atrial fibrillation and stroke prevention. Lancet Neurol 2007;6:981–93



- 120 Lip GY, Tse HF. Management of atrial fibrillation. *Lancet* 2007;370:604–18
- 121 Atrial Fibrillation Investigators. Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation. Analysis of pooled data from five randomized controlled trials. *Arch Intern Med* 1994;154:1449–57
- 122 Stroke Prevention in Atrial Fibrillation Investigators. Stroke Prevention in Atrial Fibrillation Study. Final results. *Circulation* 1991;84:527–39
- 123 Lip GY, Frison L, Halperin JL et al. Identifying patients at high risk for stroke despite anticoagulation: a comparison of contemporary stroke risk stratification schemes in an anticoagulated atrial fibrillation cohort. *Stroke* 2010;41:2731–8
- 124 Poli D, Lip GY, Antonucci E et al. Stroke risk stratification in a “real-world” elderly anticoagulated atrial fibrillation population. *J Cardiovasc Electrophysiol* 2011;22:25–30
- 125 van Staa TP, Setakis E, Di Tanna GL et al. A comparison of risk stratification schema for stroke in 79884 atrial fibrillation patients in general practice. *J Thromb Haemost* 2010;9:39–48
- 126 Dorsch MP, Lee JS, Lynch DR et al. Aspirin resistance in patients with stable coronary artery disease with and without a history of myocardial infarction. *Ann Pharmacother* 2007;41:737–41
- 127 Patel D, Moonis M. Clinical implications of aspirin resistance. *Expert Rev Cardiovasc Ther* 2007; 5:969–75
- 128 Singer DE, Albers GW, Dalen JE et al. Antithrombotic therapy in atrial fibrillation: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th edition). *Chest* 2008;133:546S–92S
- 129 Lip GY, Nieuwlaat R, Pisters R et al. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor based approach: The Euro Heart Survey on Atrial Fibrillation. *Chest* 2010;137:263–72
- 130 Pisters R, Lane DA, Nieuwlaat R et al. A novel user-friendly score (HAS-BLED) to assess 1-year risk of major bleeding in patients with atrial fibrillation: the Euro Heart Survey. *Chest* 2010;138:1093–100
- 131 Olesen JB, Lip GY, Hansen ML. Validation of risk stratification schemes for predicting stroke and thromboembolism in patients with atrial fibrillation: nationwide cohort study. *BMJ* 2011;342:d124. doi:10.1136/bmj.d124
- 132 The Task Force for the Management of Atrial Fibrillation of the European Society of Cardiology. Guidelines for the management of atrial fibrillation. *European Heart Journal* (2010) 31, 2369–2429
- 133 Iqbal MB, Taneja AK, Lip GY et al. Recent developments in atrial fibrillation. *BMJ* 2005;330:238–43
- 134 Death rates from stroke, adults aged 65 to 74, 1969 to 2006, England. [http://www.heartstats.org/temp/Figsp\\_1.1fspweb08.xls](http://www.heartstats.org/temp/Figsp_1.1fspweb08.xls). Accessed March 2009
- 135 Marini C, De Santis F, Sacco S et al. Contribution of atrial fibrillation to incidence and outcome of ischemic stroke: results from a population- based study. *Stroke* 2005;36:1115–19
- 136 Hirsh J, Dalen J, Anderson DR et al. Oral anticoagulants: mechanism of action, clinical effectiveness, and optimal therapeutic range. *Chest* 2001;119:8S–21S
- 137 Turpie AG. Warfarin replacements: Mechanisms underlying emerging agents. *Can J Cardiol* 2008;24 Suppl C:56–60C21. Aspirin for the prevention of cardiovascular disease: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2009;150:396–404
- 138 Fuster V, Ryden LE, Cannom DS et al. ACC/AHA/ESC 2006 Guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 Guidelines for the Management of Patients With Atrial Fibrillation): developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. *Circulation* 2006;114:e257–354
- 139 Ringleb PA, Bousser MG, Ford GA et al. Guidelines for management of ischaemic stroke and transient ischaemic attack 2008. *Cerebrovasc Dis* 2008;25:457–507
- 140 Lu Y, Won KA, Nelson BJ et al. Characteristics of the amiodarone- warfarin interaction during long-term follow-up. *Am J Health Syst Pharm* 2008;65:947–52
- 141 Singer DE, Albers GW, Dalen JE et al. Antithrombotic therapy in atrial fibrillation: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004;126:429S–56S
- 142 Hart RG, Pearce LA, Aguilar MI. Meta- analysis: antithrombotic therapy to prevent stroke in patients who have nonvalvular atrial fibrillation. *Ann Intern Med* 2007;146:857–67
- 143 Hart RG, Benavente O, McBride R et al. Antithrombotic therapy to prevent stroke in patients with atrial fibrillation: a meta-analysis. *Ann Intern Med* 1999;131:492–501
- 144 Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation. Analysis of pooled data from five randomized controlled trials. *Arch Intern Med* 1994;154:1449–57
- 145 Hart RG, Pearce LA, Miller VT et al. Cardioembolic vs. noncardioembolic strokes in atrial fibrillation: frequency and effect of antithrombotic agents in the stroke prevention in atrial fibrillation studies. *Cerebrovasc Dis* 2000;10:39–43
- 146 Miller VT, Pearce LA, Feinberg WM et al. Differential effect of aspirin versus warfarin on clinical stroke types in patients with atrial fibrillation. *Stroke Prevention in Atrial Fibrillation Investigators. Neurology* 1996;46: 238–40
- 147 Hylek EM. Contra: “Warfarin should be the drug of choice for thrombo- prophylaxis in elderly patients with atrial fibrillation”. Caveats regarding use of oral anticoagulant therapy among elderly patients with atrial fibrillation. *Thromb Haemost* 2008; 100:16–17
- 148 Go AS, Hylek EM, Chang Y et al. Anticoagulation therapy for stroke prevention in atrial fibrillation: how well do randomized trials translate into clinical practice? *JAMA* 2003;290:2685–92
- 149 Burton C, Isles C, Norrie J et al. The safety and adequacy of antithrombotic therapy for atrial fibrillation: a regional cohort study. *Br J Gen Pract* 2006;56:697–702
- 150 Currie CJ, Jones M, Goodfellow J et al. Evaluation of survival and ischaemic and thromboembolic event rates in patients with non-valvar atrial fibrillation in the general population when treated and untreated with warfarin. *Heart* 2006;92:196–200
- 151 Monte S, Macchia A, Pellegrini F et al. Antithrombotic treatment is strongly underused despite reducing overall mortality among high-risk elderly patients hospitalized with atrial fibrillation. *Eur Heart J* 2006;27: 2217–23
- 152 Ansell J, Hollowell J, Pengo V et al. Descriptive analysis of the process and quality of oral anticoagulation management in real-life practice in patients with chronic non-valvular atrial fibrillation: the international study of anticoagulation management (ISAM). *J Thromb Thrombolysis* 2007;23:83–91
- 153 Frykman V, Beerman B, Ryden L et al. Management of atrial fibrillation: discrepancy between guideline recommendations and actual practice exposes patients to risk for complications. *Eur Heart J* 2001;22:1954–9
- 154 Abdelhafiz AH, Wheeldon NM. Use of resources and cost implications of stroke prophylaxis with warfarin for patients with nonvalvular atrial fibrillation. *Am J Geriatr Pharmacother* 2003;1:53–60
- 155 Lightowers S, McGuire A. Cost- effectiveness of anticoagulation in nonrheumatic atrial fibrillation in the primary prevention of ischemic stroke. *Stroke* 1998;29:1827–32
- 156 Nieuwlaat R, Olsson SB, Lip GY et al. Guideline-adherent antithrombotic treatment is associated with improved outcomes compared with under- treatment in high-risk patients with atrial fibrillation. The Euro Heart Survey on Atrial Fibrillation. *Am Heart J* 2007; 153:1006–12
- 157 Baigent C, Blackwell L, Collins R et al. Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet* 2009; 373:1849–60
- 158 Szucs TD, Bramkamp M. Pharmaco- economics of anticoagulation therapy for stroke prevention in atrial fibrillation: a review. *J Thromb Haemost* 2006;4:1180–5
- 159 Appleby J, Devlin N, Parkin D. NICE’s cost effectiveness threshold. *BMJ* 2007;335:358–9
- 160 Catella-Lawson F. Vascular biology of thrombosis: platelet-vessel wall interactions and aspirin effects. *Neurology* 2001;57:S5–7
- 161 Mant J, Hobbs FD, Fletcher K et al. Warfarin versus aspirin for stroke prevention in an elderly community population with atrial fibrillation (the Birmingham Atrial Fibrillation Treatment of the Aged Study, BAFTA): a randomised controlled trial. *Lancet* 2007;370:493–503
- 162 Dorsch MP, Lee JS, Lynch DR et al. Aspirin resistance in patients with stable coronary artery disease with and without a history of myocardial infarction. *Ann Pharmacother* 2007;41:737–41
- 163 Patel D, Moonis M. Clinical implications of aspirin resistance. *Expert Rev Cardiovasc Ther* 2007; 5:969–75
- 164 Palikhe NS, Kim SH, Park HS. What do we know about the genetics of aspirin intolerance? *J Clin Pharm Ther* 2008; 33:465–72
- 165 Hughes M, Lip GY. Stroke and thromboembolism in atrial fibrillation: a systematic review of stroke risk factors, risk stratification schema and cost effectiveness data. *Thromb Haemost* 2008;99:295–304
- 166 Winter Y, Wolfram C, Schaeg M et al. Evaluation of costs and outcome in cardioembolic stroke or TIA. *J Neurol* 2009;256:954–63
- 167 van Walraven C, Hart RG, Singer DE et al. Oral anticoagulants vs aspirin in nonvalvular atrial fibrillation: an individual patient meta-analysis. *JAMA* 2002;288:2441–8
- 168 Sato H, Ishikawa K, Kitabatake A et al. Low-dose aspirin for prevention of stroke in low-risk patients with atrial fibrillation: Japan Atrial Fibrillation Stroke Trial. *Stroke* 2006;37:447–51
- 169 Gage BF, Cardinalli AB, Albers GW et al. Cost-effectiveness of warfarin and aspirin for prophylaxis of stroke in patients with nonvalvular atrial fibrillation. *JAMA* 1995;274:1839–45
- 170 Lip GY, Frison L, Grind M. Effect of hypertension on anticoagulated patients with atrial fibrillation. *Eur Heart J* 2007;28:752–9
- 171 Du X, Ninomiya T, de Galan B et al. Risks of cardiovascular events and effects of routine blood pressure lowering among patients with type 2 diabetes and atrial fibrillation: results of the ADVANCE study. *Eur Heart J* 2009;30:1128–35
- 172 Nieuwlaat R, Olsson SB, Lip GY et al. Guideline-adherent antithrombotic treatment is associated with improved outcomes compared with under- treatment in high-risk patients with atrial fibrillation. The Euro Heart Survey on Atrial Fibrillation. *Am Heart J* 2007; 153:1006–12

- 173 NICE and the National Collaborating Centre for Chronic Conditions. NICE clinical guideline 36: The management of atrial fibrillation. June 2006. <http://www.nice.org.uk/nicemedia/live/10982/30052/30052.pdf>
- 174 National institute for Health and Clinical Excellence. [http://www.nice.org.uk/about/nice/whatwedo/niceandthenhs/nice\\_and\\_the\\_nhs.jsp](http://www.nice.org.uk/about/nice/whatwedo/niceandthenhs/nice_and_the_nhs.jsp)
- 175 Lip GY, Rudolf M. The new NICE guideline on atrial fibrillation management. *Heart* 2007;93:23
- 176 The Task Force for the Management of Atrial Fibrillation of the European Society of Cardiology. Guidelines for the management of atrial fibrillation. *European Heart Journal* (2010) 31, 2369–2429
- 177 Atrial Fibrillation Investigators. Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation. Analysis of pooled data from five randomized controlled trials. *Arch Intern Med* 1994;154:1449–57
- 178 Stroke Prevention in Atrial Fibrillation Investigators. Stroke Prevention in Atrial Fibrillation Study. Final results. *Circulation* 1991;84:527–39
- 179 National Institute for Health and Clinical Excellence. Atrial fibrillation: the management of atrial fibrillation. Costing report; Implementing NICE guidance in England. July 2006. [www.nice.org.uk/nicemedia/live/10982/30061/30061.pdf](http://www.nice.org.uk/nicemedia/live/10982/30061/30061.pdf)
- 180 Bungard TJ, Ghali WA, Teo KK et al. Why do patients with atrial fibrillation not receive warfarin? *Arch Intern Med* 2000;160:41–6
- 181 Lamassa M, Di Carlo A, Pracucci G et al. Characteristics, outcome, and care of stroke associated with atrial fibrillation in Europe: data from a multicenter multinational hospital-based registry (The European Community Stroke Project). *Stroke* 2001;32:392–8
- 182 Frykman V, Beerman B, Ryden L et al. Management of atrial fibrillation: discrepancy between guideline recommendations and actual practice exposes patients to risk for complications. *Eur Heart J* 2001;22:1954–9
- 183 Deplanque D, Leys D, Parnetti L et al. Stroke prevention and atrial fibrillation: reasons leading to an inappropriate management. Main results of the SAFE II study. *Br J Clin Pharmacol* 2004;57:798–806
- 184 McBride D, Bruggenjürgen B, Roll S et al. Anticoagulation treatment for the reduction of stroke in atrial fibrillation: a cohort study to examine the gap between guidelines and routine medical practice. *J Thromb Thrombolysis* 2007;24:65–72
- 185 Friberg L, Hammar N, Ringh M et al. Stroke prophylaxis in atrial fibrillation: who gets it and who does not? Report from the Stockholm Cohort- study on Atrial Fibrillation (SCAF-study). *Eur Heart J* 2006;27: 1954–64
- 186 Nabauer M, Gerth A, Limbourg T et al. The Registry of the German Competence NETwork on Atrial Fibrillation: patient characteristics and initial management. *Europace* 2009;11:423–34
- 187 Meilitz A, Zimmermann M, Urban P et al. Atrial fibrillation management by practice cardiologists: a prospective survey on the adherence to guidelines in the real world. *Europace* 2008; 10:674–80
- 188 Bungard TJ, Ghali WA, McAlister FA et al. Physicians' perceptions of the benefits and risks of warfarin for patients with nonvalvular atrial fibrillation. *CMAJ* 2001;165:301–2
- 189 NHS Employers. Quality and Outcomes Framework guidance for GMS contract 2009/10. March 2009. [http://www.nhsemployers.org/Aboutus/Publications/Documents/QOF\\_Guidance\\_2009\\_final.pdf](http://www.nhsemployers.org/Aboutus/Publications/Documents/QOF_Guidance_2009_final.pdf)
- 190 Baruch L, Gage BF, Horrow J et al. Can patients at elevated risk of stroke treated with anticoagulants be further risk stratified? *Stroke* 2007;38: 2459–63
- 191 National Institute for Health and Clinical Excellence. Consultation on potential new indicators for the 2012/13 Quality and Outcomes Framework (QOF) 28/2/2011. <http://www.nice.org.uk/media/6BC/B1/QOFConsultation.pdf>. Accessed 8/7/2011.
- 192 Turpie AG. Warfarin replacements: Mechanisms underlying emerging agents. *Can J Cardiol* 2008;24 Suppl C:56–60C
- 193 Fuster V, Ryden LE, Cannom DS et al. ACC/AHA/ESC 2006 Guidelines for the Management of Patients with Atrial Fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 Guidelines for the Management of Patients With Atrial Fibrillation): developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. *Circulation* 2006;114:e257–e354
- 194 Singer DE, Albers GW, Dalen JE et al. Antithrombotic therapy in atrial fibrillation: American College of Chest Physicians evidence-based clinical practice guidelines (8th Edition). *Chest* 2008;133:546S–92S
- 195 Hirsh J, Dalen J, Anderson DR et al. Oral anticoagulants: mechanism of action, clinical effectiveness, and optimal therapeutic range. *Chest* 2001;119:8S–21S
- 196 Man-Son-Hing M, Laupacis A. Anticoagulant-related bleeding in older persons with atrial fibrillation: physicians' fears often unfounded. *Arch Intern Med* 2003;163:1580–6
- 197 Mant J, Hobbs FD, Fletcher K et al. Warfarin versus aspirin for stroke prevention in an elderly community population with atrial fibrillation (the Birmingham Atrial Fibrillation Treatment of the Aged Study, BAFTA): a randomised controlled trial. *Lancet* 2007;370:493–503
- 198 Anderson DR, Gardner MJ, Putnam W et al. Population-based evaluation of the management of antithrombotic therapy for atrial fibrillation. *Can J Cardiol* 2005;21:257–66
- 199 Bravata DM, Rosenbeck K, Kancir S et al. The use of warfarin in veterans with atrial fibrillation. *BMC Cardiovasc Disord* 2004;4:18
- 200 Dolan G, Smith LA, Collins S et al. Effect of setting, monitoring intensity and patient experience on anticoagulation control: a systematic review and meta-analysis of the literature. *Curr Med Res Opin* 2008;24:1459–72
- 201 Lip GY, Zarifis J, Watson RD et al. Physician variation in the management of patients with atrial fibrillation. *Heart* 1996;75:200–5
- 202 Maeda K, Sakai T, Hira K et al. Physicians' attitudes toward anti-coagulant therapy in patients with chronic atrial fibrillation. *Intern Med* 2004;43:553–60
- 203 Vasishta S, Toor F, Johansen A et al. Stroke prevention in atrial fibrillation: physicians' attitudes to anticoagulation in older people. *Arch Gerontol Geriatr* 2001;33:219–26
- 204 Hart RG, Aguilar MI. Anticoagulation in atrial fibrillation: selected controversies including optimal anticoagulation intensity, treatment of intracerebral hemorrhage. *J Thromb Thrombolysis* 2008;25:26–32
- 205 Choudhry NK, Anderson GM, Laupacis A et al. Impact of adverse events on prescribing warfarin in patients with atrial fibrillation: matched pair analysis. *BMJ* 2006;332:141–5
- 206 Tversky A, Kahneman D. Judgment under uncertainty: heuristics and biases. *Science* 1974;185:1124–31
- 207 Feinstein AR. The „chargin factor“ and qualitative decision analysis. *Arch Intern Med* 1985;145:1257–9
- 208 Devereaux PJ, Anderson DR, Gardner MJ et al. Differences between perspectives of physicians and patients on anticoagulation in patients with atrial fibrillation: observational study. *BMJ* 2001;323:1218–22
- 209 Hobbs FD, Fitzmaurice DA, Mant J et al. A randomised controlled trial and cost-effectiveness study of systematic screening (targeted and total population screening) versus routine practice for the detection of atrial fibrillation in people aged 65 and over. The SAFE study. *Health Technol Assess* 2005;9:1–74
- 210 Lip GY, Kamath S, Jafri M et al. Ethnic differences in patient perceptions of atrial fibrillation and anticoagulation therapy: the West Birmingham Atrial Fibrillation Project. *Stroke* 2002;33: 238–42
- 211 AntiCoagulation Europe. It's about time campaign. <http://www.anti-coagulationeurope.org/abouttime/survey.html>. Accessed March 2009
- 212 McCabe PJ, Schumacher K, Barnason SA. Living with atrial fibrillation: a qualitative study. *J Cardiovasc Nurs*. 2011 Jul-Aug;26(4):336-44
- 213 Dolan G, Smith LA, Collins S et al. Effect of setting, monitoring intensity and patient experience on anticoagulation control: a systematic review and meta-analysis of the literature. *Curr Med Res Opin* 2008;24:1459–72
- 214 McBride D, Bruggenjürgen B, Roll S et al. Anticoagulation treatment for the reduction of stroke in atrial fibrillation: a cohort study to examine the gap between guidelines and routine medical practice. *J Thromb Thrombolysis* 2007;24:65–72
- 215 Trummer U, Mueller U, Nowak P et al. Does physician-patient communication that aims at empowering patients improve clinical outcome? A case study. *Patient Educ Couns* 2006;61:299–306
- 216 Beyth RJ, Quinn L, Landefeld CS. A multicomponent intervention to prevent major bleeding complications in older patients receiving warfarin. A randomized, controlled trial. *Ann Intern Med* 2000;133:687–95
- 217 Heneghan C et al. Review: self testing and self management increase the benefits and reduce the harms of anticoagulant therapy. *The Lancet*, 2006; 367: 404-11
- 218 MHRA Guidance, 2004 [www.mhra.gov.uk](http://www.mhra.gov.uk)
- 219 Gardiner C et al. Patient self-testing is a reliable and acceptable alternative to laboratory INR monitoring. *British Journal of Haematology*. 2004; 128: 242-247.
- 220 Fitzmaurice DA et al. Self-management of oral anticoagulation: randomised trial. *British Medical Journal*. 2005; 331(7524): 1057.
- 221 Ansell J, Hollowell J, Pengo V et al. Descriptive analysis of the process and quality of oral anticoagulation management in real-life practice in patients with chronic non-valvular atrial fibrillation: the international study of anticoagulation management (ISAM). *J Thromb Thrombolysis* 2007;23:83–91
- 222 Lip GY, Agnelli G, Thach AA et al. Oral anticoagulation in atrial fibrillation: A pan-European patient survey. *Eur J Intern Med* 2007;18:202–8
- 223 Macik BG. The future of anticoagulation clinics. *J Thromb Thrombolysis* 2003;16:55–9
- 224 Rodgers H, Sudlow M, Dobson R et al. Warfarin anticoagulation in primary care: a regional survey of present practice and clinicians' views. *Br J Gen Pract* 1997;47:309–10
- 225 Chiquette E, Amato MG, Bussey HI. Comparison of an anticoagulation clinic with usual medical care: anticoagulation control, patient outcomes, and health care costs. *Arch Intern Med* 1998;158:1641–7
- 226 Taborski U, Wittstamm FJ, Bernardo A. Cost-effectiveness of self-managed anticoagulant therapy in Germany. *Semin Thromb Hemost* 1999;25:103–7
- 227 Murray E, Fitzmaurice D, McCahon D et al. Training for patients in a randomised controlled trial of self management of warfarin treatment. *BMJ* 2004;328:437–8
- 228 Poller L, Keown M, Ibrahim S et al. A multicentre randomised clinical endpoint study of PARMA 5 computer-assisted oral anticoagulant dosage. *Br J Haematol* 2008;143: 274–83

- 229 Poller L, Keown M, Ibrahim S et al. An international multicenter randomized study of computer-assisted oral anti-coagulant dosage vs. medical staff dosage. *J Thromb Haemost* 2008; 6:935–43
- 230 Groene O, Lombarts M, Klazinga N et al. Is patient-centredness in European hospitals related to existing quality improvement strategies? Analysis of a cross-sectional survey (MARQuIS study). *Qual Saf Health Care* 2009;18:i44–i50
- 231 Ellis S. The patient-centred care model: holistic/multiprofessional/reflective. *Br J Nurs* 1999;8:296–301
- 232 Gage BF, Cardinalli AB, Owens DK. Cost-effectiveness of preference-based antithrombotic therapy for patients with nonvalvular atrial fibrillation. *Stroke* 1998;29:1083–91
- 233 Biem HJ, Hadjistavropoulos H, Morgan D et al. Breaks in continuity of care and the rural senior transferred for medical care under regionalisation. *Int J Integr Care* 2003;3:e03
- 234 van Bommel JH, van Ginneken AM, Stam B et al. Virtual electronic patient records for shared care. *Stud Health Technol Inform* 1998;52 Pt 1: suppl 37–41
- 235 van Walraven C, Seth R, Austin PC et al. Effect of discharge summary availability during post-discharge visits on hospital readmission. *J Gen Intern Med* 2002;17:186–92
- 236 Turpie AG. New oral anticoagulants in atrial fibrillation. *Eur Heart J* 2007; 29:155–65
- 237 Turpie AG. Oral, direct factor Xa inhibitors in development for the prevention and treatment of thromboembolic diseases. *Arterioscler Thromb Vasc Biol* 2007;27:1238–47
- 238 EMEA. European Public Assessment Report for Eliquis. [http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/002148/human\\_med\\_001449.jsp&url=menus/medicines/medicines.jsp&mid=WCOB01ac058001d125#](http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/002148/human_med_001449.jsp&url=menus/medicines/medicines.jsp&mid=WCOB01ac058001d125#)
- 239 Patel MR for ROCKET AF Investigators. Rivaroxaban-once daily, oral, direct factor Xa inhibition compared with vitamin K antagonism for prevention of stroke and Embolism Trial in Atrial Fibrillation: rationale and design of the ROCKET AF study. *Am Heart J* 2010;159:340–7
- 240 Mahaffey KW, Fox KAA and ROCKET AF Executive Steering Committee. Stroke prevention using the oral direct Factor Xa inhibitor rivaroxaban compared with warfarin in patients with nonvalvular atrial fibrillation (ROCKET AF). Presented at AHA 2010. [http://sciencenews.nyamericanheart.org/sessions/late\\_breaking.shtml#rocket](http://sciencenews.nyamericanheart.org/sessions/late_breaking.shtml#rocket). Accessed March 2011
- 241 European Medicines Agency Evaluation of Medicines for HumanUse. CHMP Assessment Report for Xarelto. <http://www.emea.europa.eu/humandocs/PDFs/EPAR/xarelto/H-944-en6.pdf>. Accessed March 2009
- 242 ClinicalTrials.gov. <http://www.clinicaltrials.gov>. Accessed May 2009
- 243 Weitz JI. Randomized, parallel group, multicenter, multinational study evaluating safety of DU-176b compared with warfarin in subjects with non-valvular atrial fibrillation. 50th Annual Meeting of the American Society of Hematology, 5–9 December 2008, San Francisco: Abstract 33
- 244 European Medicines Agency. Pradaxa Annex 1 Summary of Product Characteristics. <http://www.emea.europa.eu/humandocs/PDFs/EPAR/pradaxa/emea-combined-h829en.pdf>. Accessed November 2009
- 245 Connolly SJ, Ezekowitz MD, Yusuf S et al. Dabigatran versus warfarin in patients with atrial fibrillation. *N Engl J Med* 2009;361:10.1056/nejmoa.0905561
- 246 Reddy VY, Holmes D, Shephal K et al. Safety of Percutaneous Left Atrial Appendage Closure: Results from the Watchman Left Atrial Appendage System for Embolic Protection in Patients With AF (PROTECT AF) Clinical Trial and the Continued Access Registry Circulation DOI: 10.1161/CIRCULATIONAHA.110.976449. Jan 17, 2011
- 247 ClinicalTrials.gov. RELY-ABLE long term multi-center extension of dabigatran treatment in patients with atrial fibrillation Who Completed RE-LY Trial. 2010. <http://clinicaltrials.gov/ct2/show/NCT00808067>. Accessed February 2011
- 248 Khoo CW, Tay KH, Shantsila E et al. Novel oral anticoagulants. *Int J Clin Pract* 2009;63:630–41
- 249 The ACTIVE Investigators. Effect of clopidogrel added to aspirin in patients with atrial fibrillation. *N Engl J Med* 2009;360:2066–78
- 250 Iqbal MB, Taneja AK, Lip GY et al. Recent developments in atrial fibrillation. *BMJ* 2005;330:238–43
- 251 Connolly SJ, Crijns HJ, Torp-Pedersen C et al. Analysis of stroke in ATHENA: a placebo-controlled, double-blind, parallel-arm trial to assess the efficacy of dronedarone 400 mg BID for the prevention of cardiovascular
- 252 Lee R, Kruse J, McCarthy PM. Surgery for atrial fibrillation. *Nat Rev Cardiol* 2009;6:505–13
- 253 Sick PB, Schuler G, Hauptmann KE et al. Initial worldwide experience with the WATCHMAN left atrial appendage system for stroke prevention in atrial fibrillation. *J Am Coll Cardiol* 2007;49:1490–5
- 254 AtriCure reports first human implant of the Cosgrove–Gillinov left atrial appendage occlusion system. [www.bio-medicine.org](http://www.bio-medicine.org). Accessed October 2009
- 255 Holmes DR, Reddy VY, Turi ZG et al. Percutaneous closure of the left atrial appendage versus warfarin therapy for prevention of stroke in patients with atrial fibrillation: a randomised non-inferiority trial. *Lancet* 2009;374: 534–42
- 256 McCabe DJ, Kinsella JA, Tobin WO. Left atrial appendage occlusion in non-valvular atrial fibrillation. *Lancet* 2009;374:504–6
- 257 Khoo CW, Tay KH, Shantsila E et al. Novel oral anticoagulants. *Int J Clin Pract* 2009;63:630–41
- 258 European Medicines Agency. Clopidogrel Annex 1 Summary of Product Characteristics. <http://www.emea.europa.eu/human/docs/PDFs/EPAR/clopidogrel/bms/H-974-PI-en.pdf>. Accessed March 2009
- 259 PR Newswire. Largest registry to date to provide the first-ever picture of the real global burden of atrial fibrillation (AF). Barcelona, Spain, 30 August 2009. <http://pharmalicensing.com>. Accessed October 2009
- 260 Siddique A, Butt M, Shantsila E et al. New antiplatelet drugs: beyond aspirin and clopidogrel. *Int J Clin Pract* 2009;63:776–89
- 261 Hohnloser SH, Crijns HJ, van Eickels M et al. Effect of dronedarone on cardiovascular events in atrial fibrillation. *N Engl J Med* 2009; 360:668–78
- 262 Connolly SJ, Crijns HJ, Torp-Pedersen C et al. Analysis of stroke in ATHENA: a placebo-controlled, double-blind, parallel-arm trial to assess the efficacy of dronedarone 400 mg BID for the prevention of cardiovascular hospitalization or death from any cause in patients with atrial fibrillation/atrial flutter. *Circulation* 2009;120:1174–80
- 263 Nabauer M, Gerth A, Limbourg T et al. The Registry of the German Competence NETwork on Atrial Fibrillation: patient characteristics and initial management. *Europace* 2009;11:423–34
- 264 Kakkor AK, Lip GYH, Breithardt G. The importance of real-world registries in the study of AF-related stroke. <http://www.theheart.org/documents/sitestructure/en/content/programs/1003241/transcript.pdf>. Accessed October 2009
- 265 The office of Health Economics estimating the direct costs of AF to the NHS in constituent countries of the UK
- 266 British Heart Foundation Stroke Statistics 2009
- 267 SAFE Study
- 268 Wrexham Pilot Study, 2007, Wrexham Maelor Hospital

# Atrial fibrillation—so what? Changing clinical practice

**Michelle Bennett** is British Heart Foundation Arrhythmia Nurse Practitioner, North Wales NHS Trust (East), Wrexham Maelor Hospital, Croesnewydd Road, Wrexham, LL13 7TD, UK. **Email:** michelle.bennett@new-tr.wales.nhs.uk

The arrhythmia nurse practitioner's role has many different titles, from specialist nurse to arrhythmia care co coordinator, with the different titles comes the diverse yet rewarding workload. The role of the British Heart Foundation (BHF) arrhythmia nurse is underpinned with three key elements

- ♦ Ensuring that all patients with arrhythmias receive an effective and holistic assessment and a package of care, to ensure that all the patients' medical and emotional needs are discussed.
- ♦ All those who are included in the care pathways including patients, families and carers receive education and support as needed.
- ♦ Ongoing monitoring and auditing of the arrhythmia service takes place. These elements ensure that the quality requirements of chapter 8 of the National Service Framework (NSF) for Coronary Heart Disease (CHD) (Department of Health, 2005) and the Welsh equivalent standard 5 of the Welsh NSF (Welsh Assembly Government, 2008) are met.

## Arrhythmia nurse practitioners service

In October 2006 two experienced cardiac nurses were appointed to the arrhythmia nurse practitioners (ANPs) service based in a district general hospital, one with additional experience in primary care with knowledge of the local area. These roles had been created to develop a service that would bridge the gap between primary, secondary and tertiary care for patients who suffer from arrhythmias.

The day-to-day work load now is very varied and can include answering telephone enquires from the advice line, pre-assessing patients for procedures such as elective direct current cardioversion, permanent pacemaker implants, educating other health professionals about the management of patients with arrhythmias, visiting patients, families or carers at home or in hospital to discuss arrhythmias and their management, liaising with other health professionals to improve patient care, audit, and developing patient pathways to improve access to services. Nevertheless the majority of our work load centres around the management of patients with atrial fibrillation (AF).

## Atrial fibrillation

Atrial fibrillation (AF) is the most common sustained cardiac arrhythmia, if left untreated it is a significant risk

factor for stroke and other morbidities (National Collaborating Centre for Chronic Conditions (NCC-CC), 2006). The annual risk of stroke increases by around 4-5% for patients with atrial fibrillation, however this is not relative and the risk increases with age and other co-morbidities. This risk can be reduced with appropriate and timely thromboprophylaxis (NCC-CC, 2006). The effects of AF on the patient can range from none to many side effects such as reduced quality of life, breathlessness, fatigue, palpitations, and angina. The screening for atrial fibrillation in the elderly (SAFE) study (Hobbs et al, 2005) identified that when GPs record manual pulses during routine consultations the incidence of AF diagnosis is significantly increased, ultimately leading to a reduction in the incidence of stroke.

Based on experience as a practice nurse prior to taking the post as an arrhythmia nurse practitioner, it was felt that GP surgeries and practice nurses play a vital role in the care of patients with chronic diseases and are therefore

## ABSTRACT

Atrial fibrillation (AF) is the most common sustained cardiac arrhythmia, if left untreated it is a significant risk factor for stroke and other morbidities. Approximately 12 500 strokes each year are attributable to AF and the annual cost to the NHS and personal social services budget is estimated to be around £148 million. The SAFE study identified that when GPs record manual pulses during routine consultations the incidence of AF diagnosis is significantly increased, ultimately leading to a reduction in the incidence of stroke.

On 1 May 2007 the Wrexham primary care AF model pilot was launched. Working in partnership across organizational boundaries, changes were made to existing templates in some GP surgeries. A manual pulse check was added to all chronic disease management templates, and a stroke risk stratification tool was added to AF templates to ensure patients are correctly stratified for appropriate use of a thromboprophylactic agent, which in turn would reduce the incidence of stroke. Seven new AF patients were found by opportunistic checks during the pilot and 68 patients found to be on inappropriate or no thromboprophylaxis, which prompted further review by the GPs.

## KEY WORDS

- ♦ Manual pulse checks
- ♦ Atrial fibrillation
- ♦ Screening
- ♦ Stroke risk assessment

Submitted for review 30 March 2009. Accepted for publication 23 April 2009.  
Conflict of interest: None.

ideally situated to initiate a screening service for atrial fibrillation within local primary care settings.

The hospital is based on the borders of Wales and England, serving patients from both countries, and the same region is also served by two other hospitals. To reduce inequalities in services between the different hospital that serve the local area, we chose to facilitate other health care professionals in the community to undertake screening for AF by integrating manual pulse checks into routine clinical assessments i.e. chronic disease management in GP surgeries, district nurse assessments, community matron or long-term condition nurse assessments.

### **Facilitating AF screening in GP surgeries**

The model for screening is not complex, nor ingenious, quite the opposite. It is simple, requires no funding, and can be adapted into any general practice setting that conducts chronic disease management reviews. The model aims for each individual to be assessed, to create a treatment plan that is centred on their own exclusive needs.

Before piloting the model we delivered several educational workshops for local practitioners (including GPs, practice nurses and district nurses) to reach as many health professionals in primary care as possible in order to highlight the implications of AF. Emphasis was placed on the need for manual pulse and blood pressures monitoring, as the use of automated machines can give inaccurate readings for patients with irregular pulses.

The arrhythmia nurse practitioner service worked in partnership across organizational boundaries and made changes to existing templates in some GP surgeries (who volunteered to take part). These included the addition of a manual pulse check to all chronic disease management templates, which instigates opportunistic and routine screening not only for those who are at high risk for AF but also for the wider population.

In addition, the model used a stroke risk stratification tool—CHADS<sub>2</sub> (Valentin et al, 2006)—and recorded the results using the read code 388I in an AF template to ensure patients were correctly stratified for the appropriate thromboprophylactic agent, which in turn would reduce the incidence of stroke.

The CHADS<sub>2</sub> stroke risk stratification is a clinical prediction tool used for estimating the risk of stroke in patients with nonrheumatic, or nonvalvular, AF. Points are assigned for chronic heart failure (C), hypertension (H), age 75 years or over (A), diabetes mellitus (D) and history of stroke or transient ischaemic attack (S—2 points). The higher a person's CHADS<sub>2</sub> score, the greater the risk of stroke.

An annual diary date was created to prompt annual review, which is a component missing from the clinical indicators of the quality outcome framework (QOF) for atrial fibrillation (British Medical Association/NHS Employers 2006).

Implementing the tool and annual check was supported by the development of internet-based guidelines and cardiac network guidelines that GPs can access.

### **AF primary care pilot**

The Wrexham primary care AF model pilot launched on 1st May 2007 in four general practice surgeries under the local health boards, and ran for six months with positive results and findings. The audit of the pilot evaluated the review of existing patient on the AF registers in the surgeries as well as those patients that were identified through the routine screening using the manual pulse check. Seven new AF patients were found during opportunistic checks as part of the pilot and 68 patients were found to be on inappropriate or no thromboprophylaxis, which prompted further review by the GPs.

Nearly two years since the start of the pilot the team are still reaping the benefits of the work undertaken in primary care. Although four practices were initially recruited into the pilot, the arrhythmia nurse practitioners wrote to every practice in the catchment area and asked to set up a meeting to discuss the basis of the pilot.

It seems a key reason that practices were not keen to take part in the pilot was that the pilot scheme used paper audit forms, whereas most GP surgeries operate a paperless policy. This appeared to be an obstacle that could not be overcome at the time, and the additional time needed to complete the form was seen to be problematic.

In one surgery one of the ANPs personally performed an audit of all the patients on the AF register, highlighting any patients that appeared to be on inappropriate therapy or thromboprophylaxis. This allowed her to gain a better understanding of the extent of problems that could be discovered, and appreciate the time it took to perform this audit and the impact it would have on GP and practice nurses' workload. This method of auditing patients was also used in other surgeries by GPs and practice nurses, whereas some practices chose to invite all patients on the AF register in for a review.

The ANP service now see on many referral letters that the patient was found to be in AF during a routine blood pressure check or during an annual chronic disease management review. Hopefully, this is a direct effect of the educational events the service continues to provide and also a result of those surgeries that made the changes to their chronic disease management templates.

### **Award winning service.**

In April 2008 the team's work in this area was recognized nationally with an award at the Cardiac Nursing Awards in London for 'Excellence or innovation in arrhythmia management'. This award would not have been possible if it were not for the continued support of our manager, consultants, GPs, practice nurses, cardiac lead nurses in the community and last but not least the BHF. Our idea was simple but has proved to be effective.

### **The future**

Other initiatives undertaken by the team include a collaboration with the local health board lead nurse for cardiovascular disease on developing part of a cardiology local enhanced service (LES) for GPs (DH, 2008)).

The option in the LES relating to AF included raising awareness of AF, the significance of taking a manual pulse and the importance of using the stroke risk stratification tool to identify those who needed interventions to reduce the risk of stroke. GP practices agreeing to take part in this option were asked to attend an ECG study day on arrhythmias and an AF study session provided by the two ANPs. The session included the importance of the stroke risk stratification tool and thromboprophylaxis. Following this, the practices involved were encouraged to adapt all their chronic disease management computer templates to include taking a manual pulse.

It is recognized that ongoing education is required in primary care and working closely with local health boards the ANP service aims to provide further education.

In addition, the service is in the process of organizing a 'Know your Pulse' educational event for Arrhythmia Awareness Week 2009. The event is to be hosted with the support of the occupational health department of a large local employer who has approximately 8500 permanent and contract staff. The event will start with an internal television campaign highlighting the importance of 'know your pulse' followed by site tours educating staff on how to check their own pulse and the importance of seeking medical advice if an abnormal pulse is found. Due to the enormity of the site three days have been scheduled for this event over the summer months, alongside night visits to ensure night staff benefit from this event.

## KEY POINTS

- ◆ Atrial fibrillation is the most common arrhythmia and if left untreated can lead to an increased risk of stroke
- ◆ Undetected AF cannot be treated, making screening important
- ◆ Studies have shown that when GPs record manual pulses during routine consultations the incidence of AF diagnosis is significantly increased
- ◆ By providing education and support to primary care colleagues to assist them in identifying and appropriately treating patients with AF, future complications may be reduced

- British Medical Association/NHS Employers (2006) *Revisions to the GMS Contract 2006/07*. Annex 1: Quality and Outcomes Framework guidance 2006/07 <http://tinyurl.com/d6zmlr> (accessed 27 April 2009)
- Department of Health (2005) *National Service Framework for Coronary Heart Disease. Chapter 8 Arrhythmia and Sudden Cardiac Death*. DH, London
- Department of Health (2008) Enhanced services. <http://tinyurl.com/cj38gh> (accessed 27 April 2009)
- National Collaborating Centre for Chronic Conditions (2006) *Atrial Fibrillation. National Clinical Guideline for Management in Primary and Secondary Care*. [Full version of NICE Clinical Guideline 36] Royal College of Physicians, London.
- Welsh Assembly Government (WAG) (2007) *Interim Update of Original National Service Framework. Tackling Coronary Heart disease and Arrhythmias in Wales*. WAG

Link to:

Health Technology Assessment 2005; Vol. 9: No. 40

<http://www.hta.ac.uk/fullmono/mon940.pdf>